



# Outline

- How small is small?
- What are we trying to learn?
- How we study particles?
- What does Brigitte do all day?


# Élémentaire! Mon cher Watson.

Particle physicists seek to understand what are the fundamental building blocs of Nature and how they interact to make up our Universe.





10<sup>-7</sup> m virus





virus de l'influenza

10<sup>-7</sup> m virus

10<sup>-9</sup> m

molecule





10<sup>-7</sup> m virus

10<sup>-9</sup> m molecule

10<sup>-10</sup> m atom


10<sup>-14</sup> m nucleus

10<sup>-15</sup> m proton/neutron


<10<sup>-18</sup> m

quarks






# What is matter made of?




# The ingredients of the Univers

Matter particles (6 quarks, 6 leptons)
Force carrier particles (4)



# What are we trying to understand?


- Origin of mass
- No unified description of all forces including gravity
- What is the Dark matter of the Univers?



# What are we trying to understand?

"What's the matter with antimatter?"





Matter behaves slightly differently than anti-matter (CP violation)

# How do we study elementary particles?

Use a giant "microscope": a particle accelerator






# How do we use a particle accelerator?

### **Instructions**:

- 1) Fill accelerator with a large number of particles that travel in opposite directions.
- 2) Accelerate these particles to the highest possible energy.
- 3) Bring beams of particle traveling in opposite directions into collision.
- 4) Take "pictures" of what comes out of these collisions
- 5) Analyze billions of "pictures" to study how nature works.




# Why does that work?



# **Particle collisions**

During a collision, the kinetic energy of colliding particles is converted to mass to form new massive secondary particles. This is how we can create unstable massive particles and study their properties.



# **Particle Accelerators**



### **Tevatron**

- Location: Chicago, USA
- Size: 5 km circumference
- Operation: 1992-now
- proton-antiproton collisions
- Collision energy = 2 TeV
- # collisions / s = 3 MHz
- 99.999956% speed of light
- 60,000 turns / s

### Large Hadron Collider (LHC)


- Location: Geneva, Switzerland
- Size: 27 km circumference
- Operation: 2007-future
- proton-proton collisions
- Collision energy = 14 TeV
- # collisions / s = 1 GHz
- 99.999991% speed of light
- 11,000 turns / s

# How to "see" what happens in a collision?


We use a fancy camera that takes 3D pictures: particle detector








# A "picture" of a collision




# A "picture" of a collision




## **Particle Identification**


### **Electron**




### Muon




### Quark



### **Neutrino**



### **Bottom quark**



# **Putting it all together**

**Animation** 

# **Examples from my research**


### 1- Detector construction/commissioning:

ATLAS trigger commissioning

### 2- Analysis of data


Top quark studies using the DZero experiment






# **The ATLAS Detector** 25 m 46 m


Weight = 7000 Tons








# **ATLAS Trigger System**



1 billion collisions per second

### **Trigger System**





Dr Chris Potter



Marc-Andre Dufour



Dr Cibran Santamarina

200 "photos" per second



3.2 PB data / year (4.6 Million CD = 695 year of music)


# **Data Analysis**


FERMILAB-PUB-06/475-E

### Evidence for production of single top quarks and first direct measurement of $|V_{tb}|$


The D0 Collaboration presents first evidence for the production of single top quarks at the Fermilab Tevatron  $p\bar{p}$  collider. Using a 0.9 fb<sup>-1</sup> dataset, we apply a multivariate analysis to separate signal from background and measure  $\sigma(p\bar{p} \to tb + X, tqb + X) = 4.9 \pm 1.4$  pb. The probability to measure a cross section at this value or higher in the absence of signal is 0.035%, corresponding to a 3.4 standard deviation significance. We use the cross section measurement to directly determine the CKM matrix element that describes the Wtb coupling and find 0.68 <  $|V_{tb}| \leq 1$  at 95% C.L. within the standard model.


PACS numbers: 14.65.Ha; 12.15.Ji; 13.85.Qk






# First evidence for single top quark







Gustavo Kertzscher




Dr Chris Potter



Camille Belanger-Champagner



# **Recreating the Early Universe**



13.7 billion years

**NOW** 

1 billion years

Stars form

300,000 years

Atoms form

180 seconds

Nuclei form

10<sup>-10</sup> seconds Protons/Neutrons form

 $1x10^{-12}$  s

10<sup>-34</sup> seconds?

**Quarks differentiate** 


?? Before ??

# **LHC Tunnel**



# Centre Européen de Recherche Nucléaire





## **ATLAS International Collaboration**



Alberta Carleton

### McGill

Montréal Regina Simon Fraser Toronto TRIUMF

UBC

Victoria

York

1850 Physicists & engineers150 Universities & laboratories34 Countries

